

## Gaps in Care Program Measures

### Behavioral Health

1. Antidepressant Medication Management - Acute Treatment: the percentage of members age 18 and older who were diagnosed with major depression and remained on an antidepressant for at least 12 weeks
2. Antidepressant Medication Management - Continuing Treatment: the percentage of members age 18 and older who were diagnosed with major depression and remained on an antidepressant for at least 6 months
3. Avoidance of Opioids at a High Dosage: the percentage of members who were prescribed two or more opioids on different dates that had less than 15 days of total opioid prescription coverage

### Women's Health

1. Breast Cancer Screening: the number of female members age 52-74 who've received a screening mammogram in the last 24 months
2. Cervical Cancer Screening: the number of female members age 24-64 who've received appropriate cervical cancer screening in the last 36 or 60 months
3. Chlamydia Screening in Women: the number of women ages 16-24 who are sexually active and have been screened for chlamydia in the last 12 months

### Pediatric Care

1. Immunizations for adolescents: the number of adolescents who've received at least 1 Tdap, 1 Meningococcal, and at least 2 HPV vaccines before their 13<sup>th</sup> birthday
2. Well Child Visits in the 3<sup>rd</sup>, 4<sup>th</sup>, 5<sup>th</sup>, and 6<sup>th</sup> year of life: the number of 3-6 year olds who've received at least one well child visit in the last 12 months
3. Avoidance of Antibiotic Treatment for Children with Upper Respiratory Infection: the percentage of children diagnosed with an upper respiratory infection and not dispensed an antibiotic
4. Lead Screening in Children: the number of children who at least one blood lead screening test before their 2<sup>nd</sup> birthday
5. Adolescent Well Care Visits: the number of 12-21 year olds who've received at least one well child visit in the last 12 months
6. Appropriate Testing for Children with Pharyngitis: the percentage of children age 3- 18 diagnosed with pharyngitis, dispensed an antibiotic, and received a strep test

### Diabetes Care

1. HbA1c Testing: the number of diabetic members who've received an A1c test in the last twelve months
2. Medical Attention for Nephropathy: the number of diabetic members who've either been diagnosed with Nephropathy or received nephropathy screening
3. Statin Therapy for Patients with Diabetes - Received Statin Therapy: the percentage of members age 40-75 with diabetes who were dispensed at least one statin medication during the year

## Gaps in Care Program Measures

4. **Statin Therapy for Patients with Diabetes - Statin Adherence 80%:** the percentage of members age 40-75 with diabetes who remained on a statin medication for at least 80% of the treatment period

### Respiratory Care

1. **Asthma Medication Ratio:** the number of asthmatic members who have a ratio of filled controller asthma medications to total asthma medication fills of 50% or more
2. **Pharmacotherapy Management of COPD Exacerbation Bronchodilator:** the percentage of members with COPD 40 and older who had an ED visit and were dispensed a bronchodilator
3. **Pharmacotherapy Management of COPD Exacerbation Systemic Corticosteroid:** the percentage of members with COPD 40 and older who had an ED visit and were dispensed a systemic corticosteroid

### Cardiac Care

1. **Statin Therapy for Patients with Cardiovascular Disease - Received Statin Therapy:** the percentage of male members age 21-75 and female members 40-75 with cardiovascular disease who were dispensed at least one high or moderate intensity statin medication
2. **Statin Therapy for Patients with Cardiovascular Disease - Statin Adherence 80%:** the percentage of male members age 21-75 and female members 40-75 with cardiovascular disease who remained on a high or moderate intensity statin medication for at least 80% of the treatment period